

The independence polynomial of a graph at -1

Vadim E. Levit

Department of Computer Science and Mathematics
Ariel University Center of Samaria, Ariel, Israel
levitv@ariel.ac.il

Eugen Mandrescu

Department of Computer Science
Holon Institute of Technology, Holon, Israel
eugen_m@hit.ac.il

Abstract

The *stability number* $\alpha(G)$ of the graph G is the size of a maximum stable set of G . If s_k denotes the number of stable sets of cardinality k in graph G , then $I(G; x) = s_0 + s_1x + \dots + s_\alpha x^\alpha$ is the *independence polynomial* of G [12], where $\alpha = \alpha(G)$ is the size of a maximum stable set.

In this paper we prove that $I(G; -1)$ satisfies $|I(G; -1)| \leq 2^{\nu(G)}$, where $\nu(G)$ equals the cyclomatic number of G , and the bounds are sharp.

In particular, if G is a connected well-covered graph of girth ≥ 6 , non-isomorphic to C_7 or K_2 (e.g., a well-covered tree $\neq K_2$), then $I(G; -1) = 0$.

Keywords: stable set, independence polynomial, cyclomatic number, tree, well-covered graph.

1 Introduction

Throughout this paper $G = (V, E)$ is a simple (i.e., a finite, undirected, loopless and without multiple edges) graph with vertex set $V = V(G)$ and edge set $E = E(G)$. If $X \subset V$, then $G[X]$ is the subgraph of G spanned by X . By $G - W$ we mean the subgraph $G[V - W]$, if $W \subset V(G)$. We also denote by $G - F$ the partial subgraph of G obtained by deleting the edges of F , for $F \subset E(G)$, and we write shortly $G - e$, whenever $F = \{e\}$. The *neighborhood* of a vertex $v \in V$ is the set $N_G(v) = \{w : w \in V \text{ and } vw \in E\}$, and $N_G[v] = N_G(v) \cup \{v\}$; if there is ambiguity on G , we use $N(v)$ and $N[v]$, respectively. A vertex v is *pendant* if its neighborhood contains only one vertex; an edge $e = uv$ is *pendant* if one of its endpoints is a pendant vertex. $K_n, P_n, C_n, K_{n_1, n_2, \dots, n_p}$ denote respectively, the complete graph on $n \geq 1$ vertices, the chordless path on $n \geq 1$ vertices, the chordless cycle on $n \geq 3$ vertices, and the complete multipartite graph on $n_1 + n_2 + \dots + n_p$ vertices.

The *disjoint union* of the graphs G_1, G_2 is the graph $G = G_1 \cup G_2$ having as a vertex set the disjoint union of $V(G_1), V(G_2)$, and as an edge set the disjoint union of $E(G_1), E(G_2)$. In particular, nG denotes the disjoint union of $n > 1$ copies of the graph G .

If G_1, G_2 are disjoint graphs, then their *Zykov sum* is the graph $G_1 + G_2$ with $V(G_1) \cup V(G_2)$ as a vertex set and $E(G_1) \cup E(G_2) \cup \{v_1v_2 : v_1 \in V(G_1), v_2 \in V(G_2)\}$ as an edge set.

As usual, a *tree* is an acyclic connected graph.

A *stable* set in G is a set of pairwise non-adjacent vertices. A stable set of maximum size will be referred to as a *maximum stable set* of G , and the *stability number* of G , denoted by $\alpha(G)$, is the cardinality of a maximum stable set in G .

A graph G is called *well-covered* if all its maximal stable sets are of the same cardinality [27]. If, in addition, G has no isolated vertices and its order equals $2\alpha(G)$, then G is *very well-covered* [8]. For instance, the graph $G = H \circ K_1$, obtained from H by appending a single pendant edge to each vertex of H , is very well-covered and $\alpha(G) = |V(H)|$. The following result shows that, under certain conditions, any well-covered graph has this form.

Theorem 1.1 [10] *Let G be a connected graph of girth ≥ 6 , which is isomorphic to neither C_7 nor K_1 . Then G is well-covered if and only if $G = H \circ K_1$, for some graph H of girth ≥ 6 .*

In other words, Theorem 1.1 shows that apart from K_1 and C_7 , connected well-covered graphs of girth ≥ 6 are very well-covered.

Proposition 1.2 [29] *A tree T is well-covered if and only if either $T = K_1$ or $T = H \circ K_1$ for some tree H .*

The structure of very well-covered graphs of girth at least 5 is described by the following theorem.

Theorem 1.3 [24] *Let G be a graph of girth at least 5. Then G is very well-covered if and only if $G = H \circ K_1$, for some graph H of girth ≥ 5 .*

Notice that a well-covered graph can have non-well-covered subgraphs; e.g., each subgraph of C_5 isomorphic to P_3 is not well-covered, while C_5 is well-covered.

Proposition 1.4 [7] *If G is a non-complete well-covered graph, then $G - N[v]$ is well-covered for any $v \in V(G)$.*

Let s_k be the number of stable sets in G of cardinality k , $k \in \{1, \dots, \alpha(G)\}$. The polynomial

$$I(G; x) = s_0 + s_1x + s_2x^2 + \dots + s_\alpha x^\alpha, \quad \alpha = \alpha(G),$$

is called the *independence polynomial* of G [12]. Some properties of the independence polynomial are presented in [1, 5, 14, 21, 22, 23, 24, 25]. As examples, we mention that:

$$\begin{aligned} I(G_1 \cup G_2; x) &= I(G_1; x) \cdot I(G_2; x), \\ I(G_1 + G_2; x) &= I(G_1; x) + I(G_2; x) - 1. \end{aligned}$$

The following result provides an easy recursive technique in evaluating independence polynomials of various graphs.

Proposition 1.5 [12, 14] *If $w \in V(G)$ and $uv \in E(G)$, then the following equalities hold:*

- (i) $I(G; x) = I(G - w; x) + x \cdot I(G - N[w]; x);$
- (ii) $I(G; x) = I(G - uv; x) - x^2 \cdot I(G - N(u) \cup N(v); x).$

The value of a graph polynomial at a specific point can give sometimes a very surprising information about the structure of the graph (see, for instance, [3], where the value of the so-called interlace polynomial at -1 is involved). In the case of independence polynomials, let us notice that if $I(G; x) = s_0 + s_1x + s_2x^2 + \dots + s_\alpha x^\alpha$, $\alpha = \alpha(G)$, then:

- $I(G; 1) = s_0 + s_1 + s_2 + \dots + s_\alpha = f(G)$ equals the number of stable sets of G , where $f(G)$ is known as the *Fibonacci number* of G [17, 26, 28];
- $I(G; -1) = s_0 - s_1 + s_2 - \dots + (-1)^\alpha s_\alpha = f_0(G) - f_1(G)$, where

$$f_0(G) = s_0 + s_2 + s_4 + \dots, \quad f_1(G) = s_1 + s_3 + s_5 + \dots$$

are equal to the numbers of stable sets of even size and odd size of G , respectively. $I(G; -1)$ is known as the *alternating number of independent sets* [6].

The difference $|f_0(G) - f_1(G)|$ can be indefinitely large. It is easy to check that the complete n -partite graph $K_{\alpha, \alpha, \dots, \alpha}$ is well-covered, $\alpha(K_{\alpha, \alpha, \dots, \alpha}) = \alpha$, and its independence polynomial is $I(K_{\alpha, \alpha, \dots, \alpha}; x) = n(1+x)^\alpha - (\alpha-1)$. Hence, $I(K_{\alpha, \alpha, \dots, \alpha}; -1) = 1 - \alpha$, i.e., for any negative integer k there is some connected well-covered graph G such that $I(G; -1) = k$.

Let $G_i, 1 \leq i \leq k$, be k graphs with $I(G_i; -1) = 2$ for every $i \in \{1, 2, \dots, k\}$, and $H = G_1 + G_2 + \dots + G_k$. Then,

$$I(H; x) = I(G_1; x) + I(G_2; x) + \dots + I(G_k; x) - (k-1)$$

and consequently, $I(H; -1) = 2k - (k-1) = k+1$. In other words, for any positive integer k there is some connected well-covered graph G such that $I(G; -1) = k$.

In this paper we prove that:

- $I(T; -1) \in \{-1, 0, 1\}$ for every tree T ;
- $I(G; -1) = 0$ for every connected well-covered graph G of girth ≥ 6 , non-isomorphic to C_7 or K_2 ;
- $|I(G; -1)| \leq 2^{\nu(G)}$, for every graph G , where $\nu(G)$ is its cyclomatic number.

2 $I(G; -1)$ for a graph G with at most one cycle

There are graphs G with at least one pendant vertex having $I(G; -1) \in \{-1, 0, 1\}$; see, for instance, the graphs from Figure 1, whose independence polynomials are, respectively,

$$\begin{aligned} I(G_1; x) &= 1 + 5x + 5x^2 + x^3, I(G_2; x) = (1 + 5x + 4x^2 + x^3)(1 + x), \\ I(G_3; x) &= 1 + 4x + 2x^2, I(G_4; x) = 1 + 5x + 5x^2 + 2x^3. \end{aligned}$$

Lemma 2.1 *If $u \in V(G)$ is a pendant vertex of G and $v \in N(u)$, then*

$$I(G; -1) = (-1) \cdot I(G - N[v]; -1).$$

Moreover, if G has two pendant vertices at 3 distance apart, then $I(G; -1) = 0$.




Figure 1: $I(G_1; -1) = 0 = I(G_2; -1)$, while $I(G_3; -1) = -1 = -I(G_4; -1)$.

Proof. Since $u \in V(G)$ is a pendant vertex of G and $v \in N(u)$, Proposition 1.5(i) assures that

$$\begin{aligned} I(G; x) &= I(G - v; x) + x \cdot I(G - N[v]; x) \\ &= (1 + x) \cdot I(G - \{u, v\}; x) + x \cdot I(G - N[v]; x) \end{aligned}$$

and this implies $I(G; -1) = (-1) \cdot I(G - N[v]; -1)$.

Let a, b be two pendant vertices of G with $\text{dist}(a, b) = 3$ and let $v \in N(a)$. According to Proposition 1.5(i), we get:

$$\begin{aligned} I(G; x) &= I(G - v; x) + x \cdot I(G - N[v]; x) \\ &= I(\{a\}; x) \cdot I(G - \{a, v\}; x) + x \cdot I(\{b\}; x) I(G - \{b\} \cup N[v]; x) \\ &= (1 + x) [I(G - \{a, v\}; x) + x \cdot I(G - \{b\} \cup N[v]; x)], \end{aligned}$$

which clearly implies $I(G; -1) = 0$. ■

Corollary 2.2 *If G is a well-covered graph of girth ≥ 6 and $G \neq qC_7$ for any $q \geq 1$, or G is a very well-covered graph of girth at least 5 and $G \neq qK_2$ for any $q \geq 1$, then $I(G; -1) = 0$, i.e., the number of stable sets of even size equals the number of stable sets of odd size. In particular, the assertion is true for every well-covered tree $T \neq K_2$.*

Proof. Notice that $I(K_2; x) = 1 + 2x$, and $I(C_7; x) = 1 + 7x + 14x^2 + 7x^3$. Therefore, $I(K_2; -1) = -1$, while $I(C_7; -1) = 1$.

If $G = K_1$, then $I(K_1; x) = 1 + x$ and, clearly, $I(G; -1) = 0$.

Otherwise, according to Theorems 1.1 and 1.3, it follows that $G = H \circ K_1$, for some graph H of order at least two, since $G \neq K_2$. Consequently, G has at least two pendant vertices at 3 distance apart, and by Lemma 2.1, we obtain that $I(G; -1) = 0$.

If $T \neq K_2$ is a well-covered tree, then either $T = K_1$ or T is very well-covered and its girth is greater than 5. In both cases, we get $I(T; -1) = 0$. ■

In [2] it is shown that

$$I(P_n; x) = F_{n+1}(x) \text{ and } I(C_n, x) = F_{n-1}(x) + 2x \cdot F_{n-2}(x),$$

where $F_n(x), n \geq 0$, are *Fibonacci polynomials*, i.e., the polynomials defined recursively by

$$F_0(x) = 1, F_1(x) = 1, F_n(x) = F_{n-1}(x) + x \cdot F_{n-2}(x).$$

Let us notice that $P_n, n \geq 5$, and C_n , for $n = 6$ or $n \geq 8$, are not well-covered.

Lemma 2.3 *For $n \geq 1$, the following equalities hold:*

- (i) $I(P_{3n-2}; -1) = 0$ and $I(P_{3n-1}; -1) = I(P_{3n}; -1) = (-1)^n$;
- (ii) $I(C_{3n}; -1) = 2 \cdot (-1)^n, I(C_{3n+1}; -1) = (-1)^n$ and $I(C_{3n+2}; -1) = (-1)^{n+1}$.

Proof. (i) We prove by induction on n .

For $n = 1$, we have $I(P_1; -1) = 0$ and $I(P_2; -1) = I(P_3; -1) = -1$, because

$$I(P_1; x) = 1 + x, I(P_2; x) = 1 + 2x, I(P_3; x) = 1 + 3x + x^2.$$

Assume that the assertion is true for any $k \leq 3n$. Using Proposition 1.5(i), we obtain $I(P_{k+1}; x) = I(P_k; x) + xI(P_{k-1}; x)$, which leads respectively to:

$$\begin{aligned} I(P_{3n+1}; -1) &= I(P_{3n}; -1) - I(P_{3n-1}; -1) = (-1)^n - (-1)^n = 0; \\ I(P_{3n+2}; -1) &= I(P_{3n+1}; -1) - I(P_{3n}; -1) = 0 - (-1)^n = (-1)^{n+1}; \\ I(P_{3n+3}; -1) &= I(P_{3n+2}; -1) - I(P_{3n+1}; -1) = (-1)^{n+1} - 0 = (-1)^{n+1}. \end{aligned}$$

(ii) Firstly,

$$\begin{aligned} I(C_3; x) &= 1 + 3x, & I(C_4; x) &= 1 + 4x + 2x^2, \\ I(C_5; x) &= 1 + 5x + 5x^2, & I(C_6; x) &= 1 + 6x + 9x^2 + 2x^3, \end{aligned}$$

which implies that

$$\begin{aligned} I(C_3; -1) &= 2 \cdot (-1), & I(C_{3+1}; -1) &= (-1), \\ I(C_{3+2}; -1) &= (-1)^2, & I(C_{3+2}; -1) &= 2 \cdot (-1)^2. \end{aligned}$$

Let uv be an edge of some $C_n, n \geq 7$. By Proposition 1.5(i), we deduce that

$$\begin{aligned} I(P_n; x) &= I(P_{n-1}; x) + x \cdot I(P_{n-2}; x) = \dots \\ &= (1 + 2x) \cdot I(P_{n-3}; x) + x \cdot (1 + x) \cdot I(P_{n-4}; x). \end{aligned}$$

Now, using Proposition 1.5(ii), we get

$$\begin{aligned} I(C_n; x) &= I(C_n - uv; x) - x^2 \cdot I(C_n - N(u) \cup N(v); x) \\ &= I(P_n; x) - x^2 \cdot I(P_{n-4}; x) = (1 + 2x) \cdot I(P_{n-3}; x) + x \cdot I(P_{n-4}; x). \end{aligned}$$

Hence, we obtain $I(C_n; -1) = (-1) \cdot [I(P_{n-3}; -1) + I(P_{n-4}; -1)]$. Since, by part (i),

$$I(P_{n-3}; -1) + I(P_{n-4}; -1) \in \{0 + (-1)^k, 2 \cdot (-1)^k\},$$

where k depends on $n - 3 \in \{3k - 2, 3k - 1, 3k\}$, it is easy to get that

$$I(C_{3k}; -1) = 2 \cdot (-1)^k, \quad I(C_{3k+1}; -1) = (-1)^k \text{ and } I(C_{3k+2}; -1) = (-1)^{k+1},$$

and this completes the proof. ■

Let us notice that there exist non-well-covered trees T_1, T_2, T_3 , non-isomorphic to P_n (see Figure 2), such that $I(T_1; -1) = -1, I(T_2; -1) = 0, I(T_3; -1) = 1$, because

$$\begin{aligned} I(T_1; x) &= 1 + 4x + 3x^2 + x^3, & I(T_2; x) &= 1 + 6x + 10x^2 + 6x^3 + x^4, \\ I(T_3; x) &= 1 + 7x + 15x^2 + 12x^3 + 5x^4 + x^5. \end{aligned}$$

The following theorem generalizes Lemma 2.3(i).




Figure 2: Non-well-covered trees.

Theorem 2.4 For any tree T , $I(T; -1) \in \{-1, 0, 1\}$, i.e., the number of stable sets of even size varies by at most one from the number of stable sets of odd size.

Proof. Let us observe that it is sufficient to show that $I(T; -1) \in \{-1, 0, 1\}$, because $I(T; x) = s_0 + s_1x + \dots + s_nx^n$, $\alpha = \alpha(T)$, implies

$$I(T; -1) = (s_0 + s_2 + s_4 + \dots) - (s_1 + s_3 + s_5 + \dots).$$

We prove by induction on $n = |V(T)|$.

For $n = 1$, $I(T; x) = 1 + x$ and $I(T; -1) = 0$.

For $n = 2$, $I(T; x) = 1 + 2x$ and $I(T; -1) = -1$.

For $n = 3$, $I(T; x) = 1 + 3x + x^2$ and $I(T; -1) = -1$.

For $n = 4$, only two trees are non-isomorphic, namely P_4 and $K_{1,3}$. Since

$$I(P_4; x) = 1 + 4x + 3x^2 \text{ and } I(K_{1,3}; x) = 1 + 4x + 3x^2 + x^3,$$

it follows that $I(P_4; -1) = 0$, while $I(K_{1,3}; -1) = -1$.

Finally, for $n = 5$ there are three non-isomorphic trees, namely $K_{1,4}$, T_5 and P_5 (see Figure 3). We have successively

$$I(K_{1,4}; x) = 1 + 5x + 6x^2 + 4x^3 + x^4,$$

$$I(T_5; x) = 1 + 5x + 6x^2 + 2x^3, \quad I(P_5; x) = 1 + 5x + 6x^2 + x^3,$$

which give $I(K_{1,4}; -1) = -1$, $I(T_5; -1) = 0$ and $I(P_5; -1) = 1$.




Figure 3: The three non-isomorphic trees on 5 vertices.

Let us suppose that $T = (V, E)$ is a tree with $|V| = n + 1 \geq 5$, v is a pendant vertex of T , $N(v) = \{u\}$ and $T_1, T_2, \dots, T_m$ are the trees of the forest $T - N[u]$. According to Proposition 1.5(ii), we get the following:

$$\begin{aligned} I(T; x) &= I(T - uv; x) - x^2 \cdot I(T - N(u) \cup N(v); x) = \\ &= (1 + x) \cdot I(T - v; x) - x^2 \cdot I(T - N[u]; x) = \\ &= (1 + x) \cdot I(T - v; x) - x^2 \cdot I(T_1; x) \cdot I(T_2; x) \cdot \dots \cdot I(T_m; x). \end{aligned}$$

Consequently, $I(T; -1) = -I(T_1; -1) \cdot I(T_2; -1) \cdot \dots \cdot I(T_m; -1) \in \{-1, 0, 1\}$, since every tree T_i has less than n vertices, and by the induction hypothesis, $I(T_i; -1) \in \{-1, 0, 1\}$. ■

Corollary 2.5 (i) For every tree, the number of dependent sets of even size varies by at most one from the number of dependent sets of odd size.

(ii) If T is well-covered tree and $T \neq K_2$, then the number of dependent sets of even size equals the number of dependent sets of odd size.

(iii) If T is well-covered tree and $T \neq K_2$, then the number of all stable sets and the number of all dependent sets are even.

Proof. Let p_1, p_2 , be the numbers of stable sets of the tree T , of odd and even size, respectively, and q_1, q_2 be the numbers of dependent sets in T , of odd and even size, respectively. Clearly, if $|V(T)| = n$, then $p_1 + q_1 = p_2 + q_2 = 2^{n-1}$. According to Theorem 2.4, $|p_1 - p_2| \leq 1$ and this implies $|q_1 - q_2| \leq 1$, while for a well-covered tree $T \neq K_2$, $p_1 = p_2$ and this leads to $q_1 = q_2$. Hence for well-covered trees different from K_2 both $p_1 + p_2$ and $q_1 + q_2$ are even numbers. ■

Corollary 2.6 If F is a forest, then $I(F; -1) \in \{-1, 0, 1\}$, and $I(F; -1) = 0$ whenever at least one of its components is a well-covered tree different from K_2 .

Proof. If $T_1, T_2, \dots, T_m, m \geq 1$, are the connected components of F , then

$$I(F; x) = I(T_1; x) \cdot I(T_2; x) \cdot \dots \cdot I(T_m; x),$$

because $F = T_1 \cup T_2 \cup \dots \cup T_m$. Now, the conclusions follow from Theorem 2.4 and Corollary 2.2. ■

The unicycle non-well-covered graph G_1 from Figure 4 has $I(G_1; x) = 1 + 5x + 5x^2 + x^3$ and $I(G_1; -1) = 0$.




Figure 4: $I(G_1; -1) = 0$, while $I(G_n; -1) \in \{-2, -1, 1, 2\}$, for $n \geq 4$.

Let us notice that for $n \geq 4$, the unicycle graph G_n in Figure 4 has

$$\begin{aligned} I(G_n; x) &= I(G_n - uv; x) - x^2 \cdot I(G_n - N(u) \cup N(v); x) \\ &= I(P_3; x) \cdot I(C_n; x) - x^2 \cdot (1 + x) \cdot I(P_{n-3}; x), \end{aligned}$$

and hence, by Lemma 2.3, we obtain

$$I(G_n; -1) = I(P_3; -1) \cdot I(C_n; -1) = (-1) \cdot I(C_n; -1) \in \{-2, -1, 1, 2\}.$$

The case of unicycle well-covered graphs is more specific.

Proposition 2.7 If G is a unicycle well-covered graph and $G \neq C_3$, then $I(G; -1) \in \{-1, 0, 1\}$.

Proof. Let us notice that $I(C_3; -1) = -2$, because $I(C_3; x) = 1 + 3x$.

If G is disconnected, then each cycle-free component H is a well-covered graph, which, by Corollary 2.2, contributes with $I(H; -1) \in \{-1, 0\}$ in the product that equals $I(G; -1)$. Therefore, we may assume that G is connected.

We show that the assertion is true by induction on $n = |V(G)|$.



Figure 5: All G_i , $1 \leq i \leq 4$, have C_3 as unique cycle, but only G_2, G_3 and G_4 are well-covered.

If $G \in \{C_4, C_5, C_7\}$, then Lemma 2.3 ensures that $I(G; -1) \in \{-1, 0, 1\}$.

If G contains C_3 and $G \neq C_3$, then for $n = 4$ there is no such connected graph, while for $n \in \{5, 6\}$, the only well-covered graphs are depicted in Figure 5. In each case, $I(G; -1) \in \{-1, 0, 1\}$.

Assume that the statement is true for unicycle well-covered graphs having at most n vertices, and let G be such a graph with $|V(G)| = n + 1$. Since G is connected and $G \notin \{C_3, C_4, C_5, C_7\}$, we infer that G has at least one pendant vertex. Hence Lemma 2.1 implies that $I(G; -1) = (-1) \cdot I(G - N[v]; -1)$. On the other hand, Proposition 1.4 assures that $G - N[v]$ is a well-covered graph. Let H_i , $1 \leq i \leq q$, be all the connected components of $G - N[v]$.

If $G - N[v]$ is forest, then $I(G; -1) \in \{-1, 0, 1\}$, according to Theorem 2.4.

If $G - N[v]$ is still a unicycle graph, let H_1 be the component containing the cycle. By Theorem 2.4, $I(H_2; -1) \cdots I(H_q; -1) \in \{-1, 0, 1\}$, while by induction hypothesis, $I(H_1; -1) \in \{-1, 0, 1\}$, as well. Therefore, we obtain

$$\begin{aligned} I(G; -1) &= (-1) \cdot I(G - N[v]; -1) \\ &= I(H_1; -1) \cdot I(H_2; -1) \cdots I(H_q; -1) \in \{-1, 0, 1\}, \end{aligned}$$

and this completes the proof. ■

Corollary 2.8 *If $T = (V, E)$ is well-covered tree of order ≥ 6 , then the following assertions are true:*

- (i) $I(T - v; -1) = I(T - N[v]; -1) = 0$ for any $v \in V$ such that $T - N[v] \neq qK_2$, $q \geq 1$;
- (ii) $I(T - uv; -1) = 0$ holds for any $uv \in E$;
- (iii) $I(T - N(u) \cup N(v); -1) = 0$ holds for any $uv \in E$.

Proof. (i) According to Proposition 1.5(i) and Corollary 2.2, we obtain

$$I(T; -1) = I(T - v; -1) + (-1) \cdot I(T - N[v]; -1) = 0.$$

Since $T - N[v] \neq qK_2$, $q \geq 1$, and according to Proposition 1.4, $T - N[v]$ is a forest consisting of well-covered trees, Corollary 2.6 ensures that $I(T - N[v]; -1) = 0$, which further implies that $I(T - v; -1) = 0$.

(ii) Let $uv \in E(T)$. If u is a pendant vertex, then

$$I(T - uv; x) = I(\{u\}; x) \cdot I(T - u; x) = (1 + x) \cdot I(T - u; x),$$

which assures that $I(T - uv; -1) = 0$. If none of u, v is pendant, then $T - uv$ is, by Proposition 1.2, a disjoint union of two well-covered trees T_1, T_2 , and consequently, Corollary 2.2 implies

$$I(T - uv; -1) = I(T_1; -1) \cdot I(T_2; -1) = 0,$$

because at least one of T_1, T_2 is non-isomorphic to K_2 .

(iii) By Proposition 1.5(iii), we have

$$I(T; x) = I(T - uv; x) - x^2 \cdot I(T - N(u) \cup N(v); x),$$

which yields $I(T - N(u) \cup N(v); -1) = I(T - uv; -1) - I(T; -1) = 0$. ■

Remark 2.9 *There is a non-well-covered tree T satisfying the equalities (i) - (iii) in Corollary 2.8 (see Figure 6).*




Figure 6: The tree T satisfies $I(T; -1) = I(T - v; -1) = 0$ for any vertex v .

Corollary 2.10 *For any tree T of order ≥ 2 , and for any vertex $v \in V(T)$, it is true that*

$$I(T - v; -1) \cdot I(T - N[v]; -1) \in \{0, 1\} \text{ and } I(T; -1) \cdot I(T - v; -1) \in \{0, 1\}.$$

Proof. According to Proposition 1.5(i), we get $I(T; x) = I(T - v; x) + x \cdot I(T - N[v]; x)$, which implies that:

$$I(T; -1) = I(T - v; -1) - I(T - N[v]; -1).$$

Since $T - v, T - N[v]$ are forests, Corollary 2.6 assures that

$$I(T; -1), I(T - v; -1), I(T - N[v]; -1) \in \{-1, 0, 1\}$$

and these are possible provided $I(T - v; -1), I(T - N[v]; -1) \in \{0, 1\}$.

Further, suppose $w \in V, v \notin V, F = T + v = (V \cup \{v\}, E \cup \{vw\})$. According to Proposition 1.5(i), we get

$$I(F; x) = I(F - v; x) + x \cdot I(F - N[v]; x) = I(T; x) + x \cdot I(T - w; x),$$

which implies that:

$$I(F; -1) = I(T; -1) - I(T - w; -1).$$

Since $I(F; -1) \in \{-1, 0, 1\}$, we infer that if $I(T; -1) = -1$, then $I(T - w; -1) \neq -1$, and if $I(T; -1) = 1$, then $I(T - w; -1) \neq 1$. Hence, $I(T; -1) \cdot I(T - w; -1) \in \{0, 1\}$. ■

3 $I(G; -1)$ and the cyclomatic number of G

The *cyclomatic number* $\nu(G)$ of the graph G is the dimension of the *cycle space* of G , i.e., the dimension of the linear space spanned by the edge sets of all the cycles of G . It is known that $\nu(G) = |E(G)| - |V(G)| + p$, where p is the number of connected components of G . If $e \in E(G)$ belongs to a cycle, then $G - e$ has the same number of connected components, and hence, $\nu(G - e) = |E(G) - \{e\}| - |V(G)| + p = \nu(G) - 1$.

Theorem 3.1 *For any graph G the alternating number of independent sets is bounded as follows*

$$-2^{\nu(G)} \leq I(G; -1) \leq 2^{\nu(G)},$$

where $\nu(G)$ is the cyclomatic number of G .

Proof. We prove by induction on $\nu(G)$.

If $\nu(G) = 0$, then G is a forest, and according to Corollary 2.6, we obtain

$$I(G; -1) \in \{-1, 0, 1\} = \{-2^{\nu(G)}, 0, 2^{\nu(G)}\}.$$

Assume that the assertion is true for graphs with cyclomatic number $\leq k$, and let G be a graph with $\nu(G) = k + 1$. Since $\nu(G) \geq 1$, G has at least one cycle, and if $uv \in E(G)$ belongs to some cycle of G , then $\nu(G - uv) = \nu(G) - 1 = k$. According to Proposition 1.5(ii), we get:

$$I(G; -1) = I(G - uv; -1) - (-1)^2 \cdot I(G - N(u) \cup N(v); -1),$$

which assures that

$$|I(G; -1)| \leq |I(G - uv; -1)| + |I(G - N(u) \cup N(v); -1)| \leq 2 \cdot 2^k,$$

because $\nu(G - uv) = k$ and $\nu(G - N(u) \cup N(v)) \leq k$. ■

Remark 3.2 *Let $G = qK_3$. Then, $I(G; x) = (1 + 3x)^q$ and hence, $I(G; -1) = (-2)^q = (-2)^{\nu(G)}$.*

Lemma 3.3 *Let $\{G_i : 1 \leq i \leq n\}$, $n > 1$, be a family of graphs, v a vertex belonging to none of G_i , and $H = H[v, G_1, \dots, G_n]$ be the graph obtained by joining v to some vertex $u_i \in V(G_i)$ for every $i \in \{1, 2, \dots, n\}$. Then*

$$I(H; -1) = I(G_1; -1) \cdot \dots \cdot I(G_n; -1) - I(G_1 - u_1; -1) \cdot \dots \cdot I(G_n - u_n; -1).$$

Moreover, $\nu(H) = \nu(G_1) + \dots + \nu(G_n)$ and H is connected whenever each G_i is connected.

Proof. According to Proposition 1.5(i), we have

$$\begin{aligned} I(H; x) &= I(H - v; x) + x \cdot I(H - N[v]; x) \\ &= I(G_1; x) \cdot \dots \cdot I(G_n; x) + x \cdot I(G_1 - u_1; x) \cdot \dots \cdot I(G_n - u_n; x), \end{aligned}$$

which clearly implies

$$I(H; -1) = I(G_1; -1) \cdot \dots \cdot I(G_n; -1) - I(G_1 - u_1; -1) \cdot \dots \cdot I(G_n - u_n; -1).$$

Since each cycle of H appears as a cycle in one of G_i , it follows that

$$\nu(H) = \nu(G_1) + \nu(G_2) + \dots + \nu(G_n).$$

Evidently, H is connected whenever every G_i is connected. ■

Corollary 3.4 *For any graph G , there exist H_1, H_2, H_3 such that:*

- (i) $I(H_1; -1) = (-1) \cdot I(G; -1)$ and $\nu(H_1) = \nu(G)$;
- (ii) $I(H_2; -1) = I(G; -1)$ and $\nu(H_2) = \nu(G) + 1$;
- (iii) $|I(H_3; -1)| = k \cdot I(G; -1)$ and $\nu(H_3) = \nu(G) + k - 1$.




Figure 7: The graphs H_1 and G have the same cyclomatic number.

Proof. (i) Let $H_1 = H[v, K_2, G]$ be the graph defined as in Lemma 3.3 (i.e., by joining with an edge the vertex v to one of the endpoints of K_2 , say u_1 , and to some vertex u_2 of G ; see Figure 7).

According to Lemma 3.3 we get $I(H_1; -1) = (-1) \cdot I(G; -1)$, since $I(K_2; x) = 1 + 2x$ and $I(K_2 - u_1; x) = 1 + x$.

(ii) Let $H_2 = H[v, W, G]$ be the graph depicted in Figure 8.




Figure 8: The graphs H_2 and G satisfy $\nu(H_2) = \nu(G) + 1$.

Since $I(W; x) = 1 + 5x + 5x^2$ and $I(W - u_1; x) = (1 + x)(1 + 3x)$, Lemma 3.3 implies that $I(H_2; -1) = I(G; -1)$.

(iii) Let $L_0 = K_1$ and $L_s, s \geq 1$, be the graph from Figure 9. Notice that

$$I(L_0; x) = 1, \quad I(L_1; x) = 1 + 3x,$$

while, by Proposition 1.5(ii), $I(L_s; x)$ satisfies

$$I(L_s; x) = (1 + 3x) \cdot I(L_{s-1}; x) - x^2 \cdot I(L_{s-2}; x), s \geq 2.$$

Hence, it follows that $I(L_s; -1) = (s + 1) \cdot (-1)^s, s \geq 0$.




Figure 9: The graph H_3 has $\nu(H_3) = \nu(G) + \nu(L_{k-1}) = \nu(G) + k$.

Let $H_3 = H[v, L_{k-1}, K_2, G]$ be the graph from Figure 9. Lemma 3.3 implies that

$$I(H_3; -1) = (-1) \cdot I(L_{k-1}; -1) \cdot I(G; -1) = (-1)^k \cdot k \cdot I(G; -1),$$

which clearly ensures that $|I(H_3; -1)| = k \cdot |I(G; -1)|$. ■

Lemma 3.5 For every positive integer ν , there exist connected graphs G_1, G_2 , with $\nu(G_1) = \nu(G_2) = \nu$, such that $I(G_1; -1) = \pm 2^q$ and $I(G_2; -1) = \pm(2^q - 1)$, for every $q \in \{0, 1, 2, \dots, \nu\}$.

Proof. Let us remark that, by Corollary 3.4(i), to show that there is a graph G with $I(G; -1) = p$, for some integer p , it is enough to find a graph G satisfying $|I(G; -1)| = |p|$.




Figure 10: $\nu(H_1) = \nu(H_2) = 5$, while $I(H_1; -1) = (-1)^4 \cdot 2^3$ and $I(H_2; -1) = (-1)^5 \cdot 2^4$.

Let G_1 be the graph obtained from a $P_3 = (\{u, v, w\}, \{uv, vw\})$ and joining v to the endpoints of $\nu - q$ graphs isomorphic to K_2 , while w is joined by an edge to one vertex of each of q graphs isomorphic to K_3 (see Figure 10 for two examples). It is easy to see that $\nu(G_1) = \nu$.

Notice that u is a pendant vertex of G_1 and $v \in N(u)$. Hence, using Lemma 2.1, we infer that

$$\begin{aligned} I(G_1; -1) &= (-1) \cdot I(G_1 - N[v]; -1) \\ &= (-1) \cdot I(qK_3; -1) = (-1)^{q+1} \cdot 2^q. \end{aligned}$$

Let $W_q, q \geq 2$, be the graph obtained from the star $K_{1,q}$ by adding qK_2 , such that each pendant vertex of $K_{1,q}$ is joined to the endpoints of one K_2 (see Figure 11 for W_5).




Figure 11: The graph W_5 has $I(W_5; x) = (1 + 3x)^5 + x \cdot (1 + 2x)^5$ and hence, $I(W_5; -1) = (-1)^5(2^5 - 1)$.

Then, $I(W_q; x) = (1 + 3x)^q + x \cdot (1 + 2x)^q$ and consequently, it follows that

$$I(W_q; -1) = (-2)^q - (-1)^q = (-1)^q(2^q - 1).$$

Using Corollary 3.4(ii) for $\nu - q$ times one can build a graph G_2 such that $\nu(G_2) = \nu$, while $I(G_2; -1) = I(W_q; -1) = (-1)^q(2^q - 1)$. ■

Proposition 3.6 For any positive integer $q = p_1^{\varepsilon_1} \cdot \dots \cdot p_k^{\varepsilon_k}$, where $p_1, p_2, \dots, p_k$ are prime numbers, there is a graph G such that

$$\nu(G) = \varepsilon_1(p_1 - 1) + \dots + \varepsilon_k(p_k - 1) \text{ and } I(G; -1) = q.$$

Proof. Recall that the graph L_s , depicted in Figure 9, has $I(L_s; -1) = (-1)^s \cdot (s + 1)$.

The graph G is obtained starting with a $P_3 = (\{u, v, w\}, \{uv, vw\})$ and joining w to one vertex from each of ε_j graphs isomorphic to L_{p_j-1} (see Figure 12 for an example). Firstly, it is easy to see that $\nu(G) = \varepsilon_1(p_1 - 1) + \dots + \varepsilon_k(p_k - 1)$.




Figure 12: The vertex w of G is joined to two L_1 , one L_2 and one L_3 .

Then, applying Proposition 1.5(i), we get

$$\begin{aligned} I(G; x) &= I(G - w; x) + x \cdot I(G - N[w]; x) \\ &= (1 + 2x) \cdot (I(L_{p_1-1}; x))^{\varepsilon_1} \cdot \dots \cdot (I(L_{p_k-1}; x))^{\varepsilon_k} \\ &\quad + x \cdot (1 + x) \cdot I(G - N[w] \cup \{u\}; x), \end{aligned}$$

which implies that $|I(G; -1)| = p_1^{\varepsilon_1} \cdot \dots \cdot p_k^{\varepsilon_k}$. ■

4 Conclusions

In this paper we proved that $|I(G; -1)| \leq 2^{\nu(G)}$ and we exhibit graphs satisfying $|I(G; -1)| \in \{2^{\nu(G)}, 2^{\nu(G)} - 1\}$.

Conjecture 4.1 *For every positive integer ν and each integer q such that $|q| \leq 2^\nu$, there is a connected graph G with $\nu(G) = \nu$ and $I(G; -1) = q$.*

Taking into account Corollary 3.4(i), it is enough to find a connected graph G such that $|I(G; -1)| = |q|$.

Remark 4.2 *We have the complete solution for $\nu \in \{0, 1, 2, 3\}$.*

- If $\nu(G) = 0$, then G must be a forest and for every $q \in \{-2^0, 0, 2^0\} = \{-1, 0, 1\}$, we saw above that there is a tree T such that $I(T; -1) = q$.
- If $\nu(G) = 1$, then G must be a unicycle graph and for every $q \in \{-2, -1, 0, 1, 2\}$, there is a connected graph G such that $I(G; -1) = q$.
- Let $\nu = 2$ and $q \in \{-2^2, -3, \dots, 3, 2^2\}$. According to Corollary 3.4(ii), (iii), and Lemma 3.5, it follows that there is a connected graph G such that $I(G; -1) = q$.
- Let $\nu = 3$ and $q \in \{-2^3, -7, \dots, 7, 2^3\}$. By Corollary 3.4(ii) and (iii), and Lemma 3.5, it follows that there is a connected graph G such that

$$I(G; -1) = q \in \{-2^3, -7, -6, -4, -3, -2, -1, 0, 1, 2, 3, 4, 6, 7, 2^3\}.$$

The graph G from Figure 13 has

$$\begin{aligned} I(G; x) &= I(G - v; x) + x \cdot I(G - N[v]; x) \\ &= (1 + 2x) \cdot (1 + 3x)^2 \cdot I(P_3; x) + x \cdot (1 + 2x)^3, \end{aligned}$$

which implies $I(G; -1) = 5$.




Figure 13: The graphs G and H satisfy $\nu(G) = 3$ and $\nu(H) = 4$.

- Let $\nu = 4$ and $q \in A = \{-2^4, -15, \dots, 15, 2^4\}$. By Corollary 3.4(ii) and (iii), and Lemma 3.5, it follows that there is a connected graph G such that

$$I(G; -1) = q \in A - \{-13, -11, 11, 13\}.$$

The graph H from Figure 13 has

$$\begin{aligned} I(H; x) &= I(H - v; x) + x \cdot I(H - N[v]; x) \\ &= (1 + 3x)^2 \cdot I(P_3; x) \cdot I(\triangle_2; x) + x \cdot (1 + 2x)^3 \cdot (1 + 5x + 5x^2), \end{aligned}$$

which implies $I(H; -1) = 11$. The case of $q = 13$ has not been settled yet.

References

- [1] Y. Alavi, P. J. Malde, A. J. Schwenk, P. Erdős, *The vertex independence sequence of a graph is not constrained*, Congressus Numerantium 58 (1987) 15-23.
- [2] J. L. Arocha, *Propriedades del polinomio independiente de un grafo*, Revista Ciencias Matematicas, vol.V (1984) 103-110.
- [3] P. N. Balister, B. Bollobás, J. Cutler, L. Pebody, *The interlace polynomial of graphs at -1*, European Journal of Combinatorics 23 (2002) 761-767.
- [4] M. Bläser, C. Hoffmann, *On the complexity of the interlace polynomial*, 2007, E-print, arXiv:cs.CC/0707.4565 v1.
- [5] J. I. Brown, K. Dilcher, R. J. Nowakowski, *Roots of independence polynomials of well-covered graphs*, Journal of Algebraic Combinatorics 11 (2000) 197-210.
- [6] Bousquet-Mélou, M., S. Linusson, and E. Nevo, *On the independence complex of square grids*, Journal of Algebraic Combinatorics 27 (2008) 423-450.
- [7] S. R. Campbell, *Some results on cubic well-covered graphs*, Ph. D. Thesis, 1987, Vanderbilt University, Dept. of Mathematics (1987).
- [8] O. Favaron, *Very well-covered graphs*, Discrete Mathematics 42 (1982) 177-187.
- [9] P. Fendley, K. Schoutens, H. van Eerten, *Hard squares with negative activity*, Journal of Physics A: Math. Gen. 38 (2005) 315-322.
- [10] A. Finbow, B. Hartnell and R. J. Nowakowski, *A characterization of well-covered graphs of girth 5 or greater*, Journal of Combinatorial Theory B 57 (1993) 44-68.

- [11] M. Fulmek, *Dual rook polynomials*, Discrete Mathematics **177** (1997) 67-81.
- [12] I. Gutman, F. Harary, *Generalizations of the matching polynomial*, Utilitas Mathematica **24** (1983) 97-106.
- [13] Y. O. Hamidoune, *On the number of independent k -sets in a claw-free graph*, Journal of Combinatorial Theory B **50** (1990) 241-244.
- [14] C. Hoede, X. Li, *Clique polynomials and independent set polynomials of graphs*, Discrete Mathematics **125** (1994) 219-228.
- [15] D. G. C. Horrocks, *The numbers of dependent k -sets in a graph is log concave*, Journal of Combinatorial Theory B **84** (2002) 180-185.
- [16] J. Jonsson, *Hard squares with negative activity and rhombus tilings of the plane*, The Electronic Journal of Combinatorics **13** (2006), #R67
- [17] A. Knopfmachera, R. F. Tichy, S. Wagner, V. Ziegler, *Graphs, partitions and Fibonacci numbers*, Discrete Applied Mathematics **155** (2007) 1175-1187.
- [18] V. E. Levit, E. Mandrescu, *Well-covered trees*, Congressus Numerantium **139** (1999) 101-112.
- [19] V. E. Levit, E. Mandrescu, *On well-covered trees with unimodal independence polynomials*, Congressus Numerantium **159** (2002) 193-202.
- [20] V. E. Levit, E. Mandrescu, *On unimodality of independence polynomials of some well-covered trees*, Discrete Mathematics and Theoretical Computer Science, Lecture Notes in Comput. Sci., **2731**, Springer, Berlin (2003) 237-256.
- [21] V. E. Levit, E. Mandrescu, *The independence polynomial of a graph - a survey*, Proceedings of the 1st International Conference on Algebraic Informatics, Aristotle University of Thessaloniki, Greece, 20-23 October, 2005, pp. 233-254.
- [22] V. E. Levit, E. Mandrescu, *Independence polynomials of well-covered graphs: Generic counterexamples for the unimodality conjecture*, European Journal of Combinatorics **27** (2006) 931-939.
- [23] V. E. Levit, E. Mandrescu, *Partial unimodality for independence polynomials of König-Egerváry graphs*, Congressus Numerantium **179** (2006) 109-119.
- [24] V. E. Levit, E. Mandrescu, *Some structural properties of very well-covered graphs*, Congressus Numerantium **186** (2007) 97-106.
- [25] V. E. Levit, E. Mandrescu, *On the roots of independence polynomials of almost all very well-covered graphs*, Discrete Applied Mathematics **156** (2008) 478-491.
- [26] A. S. Pedersen, P. D. Vestergaard, *The number of independent sets in unicyclic graphs*, Discrete Applied Mathematics **152** (2005) 246-256.
- [27] M. D. Plummer, *Some covering concepts in graphs*, Journal of Combinatorial Theory **8** (1970) 91-98.

- [28] H. Prodinger, R. F. Tichy, *Fibonacci numbers of graphs*, Fibonacci Quart. **20** (1982) 16-21.
- [29] G. Ravindra, *Well-covered graphs*, J. Combin. Inform. System Sci. **2** (1977) 20-21.
- [30] A. J. Schwenk, *On unimodal sequences of graphical invariants*, Journal of Combinatorial Theory B **30** (1981) 247-250.